



UNIVERSITY OF CAMBRIDGE  
Faculty of Mathematics

---

# MATHEMATICS WORKBOOK

August 22, 2023

## Introduction

The purpose of this workbook is to present a set of material that would be useful to know before starting the Tripos. It does not, of course, contain everything you should know. Rather than present a list of topics, the workbook contains questions on each topic that are supposed to be straightforward; by tackling these questions, you will see how much knowledge is expected.

Most of you will be familiar with most of the material covered here. It is still worth sketching out a solution to each question: it will be good revision and anyway you never really know that you have understood a problem until you do it. The answers are given at the end of the workbook. Note that none of the questions requires the use of a calculator.

If you find some of the material is unfamiliar, you should look it up. You will find most of it in any standard A-level (or the equivalent) text; and you may well find a helpful internet source just by googling the topic.

If a particular area is really unfamiliar to you, it would be worth doing exercises from a text book to supplement those given here. If you have difficulties with some questions, do not worry; you will have opportunities to cover the material when you get to Cambridge.

You may also find that you need to get into the swing of mathematics again after your long break (especially if you took a gap year). The best way to do this is to practice problem-solving. One source of problems is past STEP papers (available on the Cambridge Assessment STEP web site, [www.admissionstesting.org/for-test-takers/step/about-step/](http://www.admissionstesting.org/for-test-takers/step/about-step/)) which will be especially useful if you did not take STEP. If you have the STEP papers of this summer, you might like to have another go at them to get back up to speed.

Please e-mail comments or corrections to [undergrad-office@maths.cam.ac.uk](mailto:undergrad-office@maths.cam.ac.uk).

# Algebra

Although computers and even calculators are very good at algebra, all mathematicians agree that it is important to be able to do routine algebra quickly and accurately. You should be able to state elementary series expansions including binomial, sine and cosine, and ln series.

## 1. Factorization

Factorize the following polynomials:

(i)  $x^2 - 3x + 2$ ;

(ii)  $3x^3 - 3x^2 - 6x$ ;

(iii)  $x^2 - x - 1$ ;

(iv)  $x^3 - 1$ ;

(v)  $x^4 - 3x^3 - 3x^2 + 11x - 6$ .

*Notes:* In part (iii) you will need the quadratic formula to find the factors; part (iv) has one linear and one quadratic factor (or three linear factors two of which are complex); for part (v) you can use the factor theorem.

## 2. Inequalities

Find the values of  $x$  for which  $x^3 < 2x^2 + 3x$ .

## 3. Partial fractions

Express the following in partial fractions:

(i)  $\frac{2}{(x+1)(x-1)}$ ;

(ii)  $\frac{1}{x^3+1}$ ;

(iii)  $\frac{4x+1}{(x+1)^2(x-2)}$ ;

(iv)  $\frac{x^2-7}{(x-2)(x+1)}$ .

*Note:* It is best for these purposes not to use the ‘cover-up rule’; there are at least two other ways which involve elementary mathematics, whereas the cover-up rule works for more sophisticated reasons and to most users is simply a recipe (which does not always work).

## 4. Completing the square

Find the smallest value (for real  $x$  and  $y$ ) of:

(i)  $x^2 - 2x + 6$ ;

(ii)  $x^4 + 2x^2 + y^4 - 2y^2 + 3$ ;

(iii)  $\sin^2 x + 4 \sin x$ .

*Note:* Of course, you could find the smallest value by calculus, but expressing the function as a perfect square plus a remainder term is a surprisingly useful technique – for example, when integrating a function with a quadratic denominator.

## 5. Exponentials and logarithms

(i) The exponential function  $e^x$  can be defined by the series expansion

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots.$$

Use this definition to show that  $\frac{de^x}{dx} = e^x$ .

(ii) The natural log function  $\ln t$  can be defined (for  $t > 0$ ) as the inverse of the exponential function, so that  $\ln e^x = x$ . Set  $t = e^x$  and use the relationship  $\frac{dx}{dt} = 1 \bigg/ \frac{dt}{dx}$  to show that

$$\frac{d \ln t}{dt} = \frac{1}{t}.$$

(iii) Assuming that the exponential function has the property  $e^s e^t = e^{s+t}$ , prove that  $\ln(xy) = \ln x + \ln y$ .

(iv) The definition of  $a^x$  for any  $a$  is  $e^{x \ln a}$ . Prove that  $a^x a^y = a^{x+y}$  and  $a^{xb} = (ab)^x$ .

*Note:* If you have not thought of defining  $a^x$  in this way, it is worth considering how else you could give it a meaning when  $x$  is not an integer.

## 6. Binomial expansions

(i) Find the coefficient of  $x^k$  (for  $0 \leq k \leq 10$ ) in the binomial expansion of  $(2 + 3x)^{10}$ .

(ii) Use the binomial theorem to find the expansion in powers of  $x$  up to  $x^4$  of  $(1 + x + x^2)^6$ , by writing it in the form  $(1 + (x + x^2))^6$ .

(iii) Use binomial expansions to find the expansion in powers of  $x$  up to  $x^4$  of  $(1 - x^3)^6(1 - x)^{-6}$ .

(iv) Find the first four terms in the binomial expansion of  $(2 + x)^{\frac{1}{2}}$ .

## 7. Series expansions of elementary functions

Using only the series expansions  $\sin x = x - x^3/3! + x^5/5! + \cdots$ ,  $\cos x = 1 - x^2/2! + x^4/4! + \cdots$ ,  $e^x = 1 + x + x^2/2! + x^3/3! + \cdots$  and  $\ln(1 + x) = x - x^2/2 + x^3/3 + \cdots$ , find the series expansions of the following functions:

- |  |  |
|--|--|
| (i) $\tan x$ (up to the $x^5$ term);                   | (ii) $\sin x \cos x$ (up to the $x^5$ term); |
| (iii) $\frac{e^x + e^{-x}}{2}$ (up to the $x^5$ term); | (iv) $\ln(e^x)$ (up to the $x^3$ term);      |
| (v) $\frac{1 - \cos^2 x}{x^2}$ (up to the $x^2$ term). |  |

*Notes:* Do part (ii) without using a trigonometric formula, and compare your answer with the expansion for  $\sin(2x)$ . The function in part (iii) is  $\cosh x$ , of which more later. The interesting thing about the function in part (v) is that the series shows it is ‘well-behaved’ in the limit  $x \rightarrow 0$ , despite appearances.

## 8. Proof by induction

Prove by induction that the following results are valid.

$$(i) \quad a + ar + ar^2 + \cdots + ar^{n-1} = a \left( \frac{1 - r^n}{1 - r} \right).$$

$$(ii) \quad \sum_{r=1}^n r^3 = \frac{1}{4}n^2(n+1)^2.$$

*Note:* Methods of proof will be discussed in some detail in our first term course *Numbers and Sets*. Most students will have met mathematical induction at school; if you haven't, you will probably want to try out this straightforward but important method of proof.

## 9. Arithmetic and geometric progressions

(i) Find the sum of all the odd integers from 11 to 99.

(ii) Evaluate  $6 + 3 + \frac{3}{2} + \frac{3}{4} + \cdots$ .

(iii) Find  $\sin \theta + 2 \sin^3 \theta + 4 \sin^5 \theta + \cdots$ . (What ranges of values of  $\theta$  are allowed?)

(iv) Estimate roughly the approximate number of times a piece of paper has to be torn in half, placing the results of each tearing in a stack and then doing the next tearing, for the stack of paper to reach the moon.

*Note:* The dots in (ii) and (iii) indicate that the series has an infinite number of terms. For part (iii), recall that the expansion  $a(1+r+r^2+\cdots)$  only converges if  $-1 < r < 1$ . You may find the approximation  $10^3 = 2^{10}$  useful for part (iv). The distance from the Earth to the Moon is about  $4 \times 10^5$  km.

## Trigonometry

It is not necessary to learn all the various trigonometrical formulae; but you should certainly know what is available. The double-angle formulae, such as  $\tan 2x = 2 \tan x / (1 - \tan^2 x)$  are worth knowing, as are the basic formulae for  $\sin(A \pm B)$ ,  $\cos(A \pm B)$  and  $\tan(A \pm B)$ , the Pythagoras-type identities, such as  $\sec^2 x = 1 + \tan^2 x$ , and a few special values, such as  $\sin \pi/4 = 1/\sqrt{2}$ , that can be deduced from right-angled triangles with sides  $(1, 1, \sqrt{2})$  or  $(1, \sqrt{3}, 2)$ .

## 10. Basic identities

Starting from the identity  $\sin(A + B) = \sin A \cos B + \cos A \sin B$ , use the basic properties of the trigonometric functions (such as  $\sin(-A) = -\sin A$ ) to prove the following:

$$(i) \quad \sin(A - B) = \sin A \cos B - \cos A \sin B;$$

$$(ii) \quad \cos(A + B) = \cos A \cos B - \sin A \sin B;$$

$$(iii) \quad \tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B};$$

$$(iv) \quad \sin C + \sin D = 2 \sin \frac{1}{2}(C + D) \cos \frac{1}{2}(C - D);$$

$$(v) \quad \tan^{-1} a + \tan^{-1} b = \tan^{-1} \left( \frac{a + b}{1 - ab} \right) + n\pi.$$

*Notes:* For part (ii), recall that  $\cos A = \sin(\pi/2 - A)$ . You can use part (iii) to help with part (v). What exactly does part (v) mean?

## 11. Trigonometric equations

(i) Solve the following equations:

$$(a) \quad \sin\left(x + \frac{\pi}{6}\right) + \sin\left(x - \frac{\pi}{6}\right) = \frac{\sqrt{3}}{2}; \quad (b) \quad \cos x + \cos 2x + \cos 3x = 0.$$

(ii) Write down the value of  $\cot(\pi/6)$  and use a double angle formula to show that  $\cot(\pi/12)$  satisfies the equation  $c^2 - 2\sqrt{3}c - 1 = 0$ . Deduce that  $\cot(\pi/12) = 2 + \sqrt{3}$ .

## 12. Trigonometric identities using Pythagoras

Prove the following identities:

$$(i) \quad \tan \theta + \cot \theta = \sec \theta \operatorname{cosec} \theta; \quad (ii) \quad (\cot \theta + \operatorname{cosec} \theta)^2 = \frac{1 + \cos \theta}{1 - \cos \theta};$$
$$(iii) \quad \cos \theta = \frac{1 - t^2}{1 + t^2}; \quad \sin \theta = \frac{2t}{1 + t^2}; \quad \tan \theta = \frac{2t}{1 - t^2}, \quad \text{where } t = \tan \frac{1}{2}\theta.$$

## Complex Numbers

A complex number  $z$  can be written as  $x + iy$ , where  $x$  is the real part,  $y$  is the imaginary part and  $i^2 = -1$ . The modulus of  $z$  (written  $|z|$  or  $r$ ) is  $\sqrt{(x^2 + y^2)}$  and the argument (written  $\arg z$  or  $\theta$ ) is defined by  $x = r \cos \theta$ ,  $y = r \sin \theta$  and  $-\pi < \theta \leq \pi$ . The complex conjugate of  $z$  (written  $z^*$ ) is  $x - iy$ . The inverse,  $z^{-1}$ , of  $z$  is the complex number that satisfies  $z^{-1}z = 1$  (for  $z \neq 0$ ).

## 13. Algebra of complex numbers

Use the definitions above with  $z_1 = x_1 + iy_1$  and  $z_2 = x_2 + iy_2$  to show:

$$(i) \quad z_1 z_2 = (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + y_1 x_2); \quad (ii) \quad |z|^2 = z z^*;$$
$$(iii) \quad z^{-1} = \frac{z^*}{|z|^2}; \quad (iv) \quad |z_1 z_2| = |z_1| |z_2|;$$
$$(v) \quad \arg(z_1 z_2) = \arg z_1 + \arg z_2 \quad (\text{assume that } 0 < \arg z_1 \leq \pi/4 \text{ and } 0 < \arg z_2 \leq \pi/4).$$

Give a sketch of the  $x$ - $y$  plane (called also the complex plane or the Argand diagram) showing the points representing the complex numbers  $z_1 = 1 - i$ ,  $z_2 = -\sqrt{3} + i$ . Verify results (iii), (iv) and (v) for these numbers.

## 14. De Moivre's theorem

De Moivre's theorem is not in the FP1 syllabus for most examination boards, but it is so important that you should get to know it if you haven't already met it.

(i) Show by means of series expansions that

$$\cos \theta + i \sin \theta = e^{i\theta}$$

and deduce that  $\cos \theta - i \sin \theta = e^{-i\theta}$  and that  $z = r e^{i\theta}$ .

Deduce also that  $\cos \theta = \frac{1}{2}(e^{i\theta} + e^{-i\theta})$  and  $\sin \theta = \frac{1}{2i}(e^{i\theta} - e^{-i\theta})$ .

(ii) Use the above result to show that  $r e^{i\theta} = 1$  (for real  $\theta$  and  $r > 0$ ) if and only if  $r = 1$  and  $\theta = 2n\pi$  for some integer  $n$ .

(iii) Use part (ii) to find the three distinct roots of the equation  $z^3 = 1$ . Draw them on the complex plane and convert them from modulus-argument form to real-imaginary form.



## 18. Basic definitions

- (i) Parabola. The point  $(x, y)$  has the property that its distance from the point  $(a, 0)$  is equal to its distance from the line  $x = -a$ . Sketch the locus of the point using this information. Show that  $y^2 = 4ax$ .
- (ii) Ellipse and hyperbola. The point  $(x, y)$  has the property that its distance from the point  $(ae, 0)$  is  $e$  times its distance from the line  $x = ae^{-1}$ . Sketch the locus of the point for  $0 < e < 1$  and for  $e > 1$ . Show that  $x^2/a^2 + y^2/b^2 = 1$  if  $0 < e < 1$  and that  $x^2/a^2 - y^2/b^2 = 1$  if  $e > 1$ , where  $b = a\sqrt{|1 - e^2|}$ .
- (iii) Rectangular hyperbola. Show that the hyperbola  $x^2/a^2 - y^2/b^2 = 1$  becomes the rectangular hyperbola  $XY = 1$  in the new coordinates given by  $x = a(X + Y)/2$ ,  $y = b(X - Y)/2$ .

## 19. Parametric equations

Show that the curves with parametric form  $(x, y) = (at^2, 2at)$ ,  $(a \cos \theta, b \sin \theta)$  and  $(a \cosh \theta, b \sinh \theta)$  are conic sections.

## 20. Polar coordinates

By converting to cartesian coordinates, show that:

- (i)  $r \cos(\theta - \alpha) = c$  describes a straight line;
- (ii)  $r = \ell \cos \theta$  describes a circle;
- (iii)  $r^2 \cos(\theta + \alpha) \cos(\theta - \alpha) = c^2$  describes a hyperbola;
- (iv)  $r^{-1} = k \cos \theta + m$  can describe any conic section, the type depending on the values of  $k$  and  $m$ .

*Notes:* Here,  $c$ ,  $k$ ,  $\ell$ ,  $m$  and  $\alpha$  are constants. Special cases, such as  $\alpha = 0$  in part (iii), can be ignored. In each case, you should give a careful description ( $x$  and  $y$  intercepts,  $e$ , etc) of the curves.

# Matrices and Vectors

Matrices and vectors arise in all branches of mathematics and form an indispensable part of a mathematician's toolkit. You will have met them in FP1 (or the equivalent if you did not take A-levels) and should be familiar with the basic properties. It is also worth understanding the basic geometric uses of vectors.

We consider here only  $2 \times 2$  matrices

## 21. Matrix multiplication

Let  $\mathbf{A} = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix}$ ,  $\mathbf{B} = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$ ,  $\mathbf{C} = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$  and  $\mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ .

- (i) Show that  $\mathbf{AB} \neq \mathbf{BA}$ . (This shows that matrix multiplication is not commutative.)
- (ii) Show that  $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$ . (This illustrates that matrix multiplication is associative.)
- (iii) Without using any formulae, find a matrix  $\mathbf{D} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  such that  $\mathbf{AD} = \mathbf{I}$ .
- (iv) Show that  $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$ .
- (v) Show that  $\det \mathbf{AB} = \det \mathbf{A} \det \mathbf{B}$ .

*Notes:* The symbol  $^T$  denotes the transpose of the matrix, i.e. the matrix obtained by exchanging rows and columns:  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}^T = \begin{pmatrix} a & c \\ b & d \end{pmatrix}$ . The determinant of  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  is  $ad - bc$ .



## 22. Position vectors

Show that the points with position vectors

$$\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 4 \\ 3 \\ -2 \end{pmatrix},$$

lie on a straight line and give the equation of the line in the two forms

$$(i) \quad \mathbf{r} = \mathbf{a} + \lambda \mathbf{b}, \quad (ii) \quad \frac{x - x_0}{c} = \frac{y - y_0}{d} = \frac{z - z_0}{e}.$$

## 23. Scalar products

The three vectors  $\mathbf{A}$ ,  $\mathbf{B}$  and  $\mathbf{C}$  are defined by

$$\mathbf{A} = (1, 3, 4), \quad \mathbf{B} = (2, 1, 3), \quad \mathbf{C} = (3, 3, 2).$$

- (i) Order the vectors by magnitude.
- (ii) Use the scalar product to find the angles between the pairs of vectors (a)  $\mathbf{A}$  and  $\mathbf{B}$ , (b)  $\mathbf{B}$  and  $\mathbf{C}$ , leaving your answer in the form of an inverse cosine.
- (iii) Find the lengths of the projections of the vectors (a)  $\mathbf{A}$  onto  $\mathbf{B}$ , (b)  $\mathbf{B}$  onto  $\mathbf{A}$ .
- (iv) Find an equation of the plane, in the form  $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b} + \mu \mathbf{c}$ , through the points with position vectors  $\mathbf{A}$ ,  $\mathbf{B}$  and  $\mathbf{C}$ . Show that the normal to this plane is  $(1, 0, 1)$  and find an equation of the plane in the form  $\mathbf{r} \cdot \mathbf{n} = p$ , where  $\mathbf{n}$  is a unit vector.

*Note:* The projection of  $\mathbf{A}$  onto  $\mathbf{B}$  is the component of  $\mathbf{A}$  in the direction of the vector  $\mathbf{B}$ .

## Differentiation

Differentiation of standard functions, products, quotients, implicit function expressions, and functions of a function (using the chain rule) should be routine.

### 24. Direct differentiation

Differentiate  $y(x)$  with respect to  $x$  in the following cases:

$$\begin{array}{ll} (i) & y = \ln(x + \sqrt{1 + x^2}); \\ (ii) & y = a^x; \\ (iii) & y = x^x; \\ (iv) & y = \sin^{-1} \frac{x}{\sqrt{1 + x^2}}. \end{array}$$

*Notes:* Simplify your answer to part (i). For (ii), see the definition in question 5(iv). Can you see why the answer to (iv) is surprisingly simple?

### 25. Parametric differentiation

Show that if  $x = a \cos \theta$ ,  $y = b \sin \theta$  then  $\frac{d^2y}{dx^2} < 0$  for  $y > 0$ .

### 26. Stationary points

Find the stationary points of the function

$$f(x) = \frac{x}{x^2 + a^2},$$

where  $a > 0$ , classifying them as either maximum or minimum. Sketch the curve (without using a calculator).

## Integration

You need to be able to recognise standard integrals (without having to leaf through a formula book) and evaluate them (referring to your formula book, if necessary). You need to be familiar with the techniques of integration by parts and by substitution.

### 27. Indefinite integrals

Calculate the following integrals:

(i) $\int a^x dx;$	(ii) $\int \frac{1}{x^2 - 2x + 6} dx;$
(iii) $\int e^{ax} \cos(bx) dx;$	(iv) $\int e^{ax} e^{ibx} dx;$
(v) $\int \frac{1}{1 - x^3} dx;$	(vi) $\int \operatorname{cosec} x dx;$
(vii) $\int \sec x dx;$	(viii) $\int \frac{1}{\sqrt{(c^2 + m^2 y^2)}} dy;$
(ix) $\int \tan^{-1} x dx;$	(x) $\int x^3 e^{x^2} dx.$

*Notes:* For part (i), see question 5(iv). For part (ii), see question 4(i). Note that you can obtain (iii) from (iv) by taking the real part. Use partial fractions for part (v). For (vi) and (vii), use the substitution  $t = \tan(x/2)$  rather than the trick of multiplying top and bottom by e.g.  $\sec x + \tan x$ . Try also deriving (vii) from (vi) by means of the substitution  $y = \pi/2 - x$ . For (viii), either substitute  $my = c \sinh \theta$  or, if you are not keen on hyperbolic functions, see question 24(i). Use integration by parts for (ix) and (x).

## Differential Equations

There is a course on differential equations in the first term for which very little knowledge is assumed.

### 28. First-order equations

Find the general solution (i.e. with a constant of integration) of the following equations.

(i) $y \frac{dy}{dx} = x;$	(ii) $\frac{dy}{dx} = my;$
(iii) $x \frac{dy}{dx} = y + 1;$	(iv) $\frac{dy}{dx} - y \tan x = 1;$
(v) $\frac{dy}{dx} = \sqrt{c^2 - k^2 y^2};$	(vi) $z \frac{dz}{dy} + k^2 y = 0.$

*Note:* For part (iv), find a function ('integrating factor')  $w(x)$  such that the equation can be written  $\frac{d(wy)}{dx} = w$ .

## 29. Second-order equations

(i) Show, by substituting  $y = e^{mx}$  into the equation, that there are two values of  $m$  for which  $e^{mx}$  satisfies

$$a \frac{d^2y}{dx^2} + b \frac{dy}{dx} + cy = 0, \quad (*)$$

where  $a$ ,  $b$  and  $c$  are constants.

(ii) Show (by substitution) that if both  $y_1$  and  $y_2$  satisfy the equation (\*), then so also does the function  $y$  defined by  $y = Ay_1 + By_2$ , where  $A$  and  $B$  are constants.

(iii) Find the two values of  $m$  for which  $e^{mx}$  satisfies the equation

$$\frac{d^2y}{dx^2} + 3 \frac{dy}{dx} + 2y = 0$$

and write down a solution that contains two arbitrary constants.

You have just solved the above equation ( $y'' + 3y' + 2y = 0$ ) by guessing solutions. Instead, let  $z = y' + 2y$ . Then show that  $z' + z = 0$ , solve for  $z$  and then solve the resulting first-order differential equation of  $y$ , thereby proving that the solution you found above is in fact the most general solution.

(iv) Show by substitution that  $e^{px} \sin qx$  satisfies the equation

$$\frac{d^2y}{dx^2} - 2p \frac{dy}{dx} + (p^2 + q^2)y = 0.$$

Find the general solution by the method outlined in the second paragraph of (iii) above.

## 30. Simple harmonic motion

(i) Show that the equation

$$\frac{d^2y}{dx^2} + k^2y = 0 \quad (y \text{ real})$$

can be written as  $z \frac{dz}{dy} + k^2y = 0$ , where  $z = \frac{dy}{dx}$ . Show that  $z = \pm \sqrt{c^2 - k^2y^2}$ , where  $c$  is a constant of integration, (compare questions 28(v) and 28(vi)) and hence show that

$$y = R \sin k(x - x_0),$$

where  $R$  and  $x_0$  are constants. ( $R = c/k$  and  $x_0$  is a new constant of integration.)

(ii) Repeat the steps of part (i) on the equation

$$\frac{d^2y}{dx^2} - k^2y = 0$$

to obtain

$$y = R \sinh k(x - x_0).$$

In this case, is it the general solution?

## Answers

1. (i)  $(x-2)(x-1)$  (ii)  $3x(x-2)(x+1)$   
 (iii)  $(x - \frac{1}{2}(1 - \sqrt{5}))(x - \frac{1}{2}(1 + \sqrt{5}))$  (iv)  $(x-1)(x^2+x+1)$   
 (v)  $(x-3)(x-1)^2(x+2)$ .
2.  $x < -1$  or  $0 < x < 3$ .
3. (i)  $\frac{1}{x-1} - \frac{1}{x+1}$  (ii)  $\frac{1}{3} \left( \frac{1}{x+1} - \frac{x-2}{x^2-x+1} \right)$   
 (iii)  $\frac{1}{(x+1)^2} - \frac{1}{x+1} + \frac{1}{x-2}$  (iv)  $1 - \frac{1}{x-2} + \frac{2}{x+1}$ .
4. (i) 5; (ii) 2; (iii) -3 (smallest when  $\sin x = -1$ ).
6. (i)  $\frac{3^k 2^{10-k} 10!}{k!(10-k)!}$  (ii)  $1 + 6x + 21x^2 + 50x^3 + 90x^4$   
 (iii) Same as (ii) (iv)  $\sqrt{2} \left( 1 + \frac{1}{4}x - \frac{1}{32}x^2 + \frac{1}{128}x^3 \right)$ .
7. (i)  $x + \frac{1}{3}x^3 + \frac{2}{15}x^5$  (ii)  $x - \frac{2}{3}x^3 + \frac{2}{15}x^5$   
 (iii)  $1 + \frac{1}{2}x^2 + \frac{1}{24}x^4$  (iv)  $x$   
 (v)  $1 - \frac{1}{3}x^2$ .
9. (i) 2475 (ii) 12  
 (iii)  $\frac{\sin \theta}{\cos 2\theta}$  ( $-\pi/4 + n\pi < \theta < \pi/4 + n\pi$ ) (iv) about 42 times.
11. (ia)  $n\pi + (-1)^n \pi/6$  (ib)  $n\pi/2 + \pi/4, 2n\pi \pm 2\pi/3$ .
13.  $|z_1| = \sqrt{2}, \arg z_1 = -\pi/4, |z_2| = 2, \arg z_2 = 5\pi/6$ .
14. (iii) 1,  $e^{2i\pi/3}, e^{-2i\pi/3}$  or 1,  $(-1 \pm i\sqrt{3})/2$ .
21. (iii)  $\begin{pmatrix} 3 & -2 \\ -1 & 1 \end{pmatrix}$ .
22. (i)  $\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$  (ii)  $x-1 = y = 1-z$
23. (i)  $|\mathbf{A}| > |\mathbf{C}| > |\mathbf{B}|$ ; (ii)  $\cos^{-1} 17/(2\sqrt{91}), \cos^{-1} 15/(2\sqrt{77}),$  (iii)  $17/\sqrt{14}, 17/\sqrt{26}$   
 (iv)  $\mathbf{r} = \begin{pmatrix} 1 \\ 3 \\ 4 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -2 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ 0 \\ -2 \end{pmatrix}, \mathbf{r} \cdot \begin{pmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{pmatrix} = 5/\sqrt{2}$ .
24. (i)  $\frac{1}{\sqrt{x^2+1}}$  (ii)  $a^x \ln a$   
 (iii)  $x^x(1 + \ln x)$  (iv)  $\frac{1}{1+x^2} \quad (\sin^{-1} \frac{x}{\sqrt{x^2+1}} = \tan^{-1} x)$ .

26. Maximum at  $\left(a, \frac{1}{2a}\right)$ , minimum at  $\left(-a, -\frac{1}{2a}\right)$ .

27. (i)  $\frac{1}{\ln a} a^x + C$  (ii)  $\frac{1}{\sqrt{5}} \tan^{-1} \frac{x-1}{\sqrt{5}} + C$

(iii)  $\frac{1}{a^2 + b^2} (a \cos bx + b \sin bx) e^{ax} + C$  (iv)  $\frac{1}{a + ib} e^{(a+ib)x} + C$

(v)  $\frac{1}{6} \ln \frac{x^2 + x + 1}{(x-1)^2} + \frac{1}{\sqrt{3}} \tan^{-1} \left( \frac{2x+1}{\sqrt{3}} \right) + C$

(vi)  $\ln \tan(x/2) + C$  (vii)  $\ln \left( \frac{1 + \tan(x/2)}{1 - \tan(x/2)} \right) + C$

(viii)  $m^{-1} \ln \left( (my + \sqrt{m^2 y^2 + c^2})/c \right) + C$  or  $m^{-1} \sinh^{-1}(my/c) + C$

(ix)  $x \tan^{-1} x - \frac{1}{2} \ln(1 + x^2) + C$  (x)  $\frac{1}{2} (x^2 - 1) e^{x^2} + C$

28. (i)  $y^2 = x^2 + C$  (ii)  $y = C e^{mx}$

(iii)  $y = Cx - 1$  (iv)  $y = C \sec x + \tan x$

(v)  $ky = c \sin k(x - x_0)$  (vi)  $z^2 + k^2 y^2 = C$

29. (i) Solution of  $am^2 + bm + c = 0$  (iii)  $m = -1$  and  $m = -2$ ;  $y = Ae^{-x} + Be^{-2x}$

(iv) Set  $z = y' - (p + iq)y$ .